



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

the float. Since the whole surface of the sieve tubes in this portion of the plant is thus exposed to the gas contained in this float, it would seem possible that considerable oxidation of foods is carried on in this internal atmosphere. The gas in this float is shown to contain a little larger percentage of oxygen than air. It may possibly be worth while to consider the presence of carbon monoxide in plants in connection with the wide distribution of oxidases in plant tissue and the possible mechanism of their reaction.<sup>6</sup> LANGDON'S thorough demonstration of the presence of carbon monoxide in this cavity is a very important piece of work, and great interest attaches to the possible relation of this gas to the metabolism of the plant.—G. B. RIGG.

**Quantitative characters in beans.**—By means of a statistical study of pole and bush beans, EMERSON<sup>7</sup> has analyzed the characters causing height variation in *Phaseolus vulgaris*. They are 3 in number and apparently segregate independently after crossing. First is the manner of growth, which is either "determinate" (bush type) or "indeterminate" (pole type), with the indeterminate habit completely dominant in the  $F_1$  generation, and showing the typical 3:1 splitting in the  $F_2$  generation. Such behavior he interprets as the result of a single pair of freely segregating factors behaving in a Mendelian fashion.

TSCHERMAK, using the hybrids *Phaseolus vulgaris* × *P. multiflorus* and the reciprocal, found anomalous splitting in the  $F_2$ , since some of the "short" segregates produced "talls" in succeeding generations. He makes no mention of habit of growth, and merely classifies the progenies as "talls" and "shorts." The results of TSCHERMAK need not be compared with EMERSON'S, however, because in the former case the hybrids are interspecific, and in the latter intervarietal (intraspecific).

The second character operative in determining height is number of internodes. The presence of this character was deduced from the fact that different varieties of both pole and bush beans differed in the number of internodes produced when grown under the same conditions. The question then arose as to whether this tendency to produce few or many internodes could be inherited independently of habit of growth. Suitable crosses were made and the results seemed to answer the question in the affirmative, although the evidence is admittedly incomplete. The factors determining this difference could not be shown to be perfectly dominant, but apparent segregation followed hybridization. This segregation was attended in the  $F_2$  generation by a range of variability exceeding that of the 2 parents. EMERSON interprets this result as due to the action of multiple segregating factors.

The third character involved in height is length of internode. The modification of this character by habit of growth made its behavior difficult to study.

<sup>6</sup> REED, G. B., BOT. GAZ. 62:53-64. 1916.

<sup>7</sup> EMERSON, R. A., A genetic study of plant height in *Phaseolus vulgaris*. Research Bull. no. 7. Nebr. Agric. Exp. Sta. pp. 73. 1916.

In order to have a standard of comparison between pole and bush bean types, the first 5 internodes were measured and the means computed. For comparison between different varieties of pole beans the mean of the first 5 internodes was used. It was thought that the actual internode length found for some of the bush varieties might not be representative of the potential length which would have been attained by the upper internodes had not the production of a terminal inflorescence hindered further growth. To test this supposition crosses were made between a bush bean with long internodes and a pole bean with short internodes. The resulting hybrid showed an intermediate development in the  $F_1$  and a wide range of variation in the  $F_2$  generation. Bush beans with shorter internodes and pole beans with longer internodes than the parent types exhibited were obtained. Here again the variations were attributed to the action of multiple, non-dominant, independently segregating factors.

In conclusion, the author points out that the results of other investigators tend to show that quantitative characters in plants are inherited in two ways: (a) they are due to the action of a single Mendelian pair of factors showing complete dominance in the  $F_1$  and a 3:1 ratio in the  $F_2$  generation; (b) they exhibit an intermediate development in the  $F_1$  and a wide range of variation in the  $F_2$  generation. In class (a) belongs the determinate as opposed to the indeterminate habit of growth. Characters such as length and number of internodes fall into class (b). Such characters as those of class (b) have been interpreted in 2 ways. EMERSON, TSCHERMAK, EAST, and others attribute them to the interaction of many independently segregating factors, a theory in accord with the multiple factor hypothesis of NILSSON-EHLE. CASTLE, however, has interpreted such behavior as due, in some cases, to the modification of a unit factor through hybridization. In the case of the bean crosses, EMERSON implies that the factor involved would be that which determines habit of growth. After discussing this latter hypothesis and the assumptions its adoption would necessitate, he rejects it in favor of the multiple factor hypothesis.

Since, therefore, the characters involved in producing an effect seem to behave in different manners in inheritance, the author explains the variation in height following hybridization between pole and bush beans as due to the modification of the expression of a unit factor by the presence or absence of a number of factors producing other effects (as, for example, the effect of the determinate habit of growth on the potential length of internodes and on the number of internodes, etc.). However, the author disavows any intention of maintaining that this is the only possible explanation, and suggests that it may have to be modified to suit the results of further selection and hybridization experiments.—WILBUR BROTHERTON.

**Philippine forests.**—Our knowledge of the economic importance and the environmental conditions of some tropical forests has been advanced by a